

OCR

Oxford Cambridge and RSA

Monday 19 October 2020 – Morning

A Level Chemistry A

H432/03 Unified chemistry

Time allowed: 1 hour 30 minutes

You must have:

- the Data Sheet for Chemistry A

You can use:

- a scientific or graphical calculator
- an HB pencil



Please write clearly in black ink. Do not write in the barcodes.

Centre number

--	--	--	--	--	--

Candidate number

--	--	--	--	--

First name(s)

Last name

INSTRUCTIONS

- Use black ink. You can use an HB pencil, but only for graphs and diagrams.
- Write your answer to each question in the space provided. If you need extra space use the lined pages at the end of this booklet. The question numbers must be clearly shown.
- Answer all the questions.
- Where appropriate, your answer should be supported with working. Marks might be given for using a correct method, even if your answer is wrong.

INFORMATION

- The total mark for this paper is 70.
- The marks for each question are shown in brackets [].
- Quality of extended response will be assessed in questions marked with an asterisk (*).
- This document has 20 pages.


ADVICE

- Read each question carefully before you start your answer.

1 This question is about organic chemistry.


(a) This part is about two practical techniques used in organic preparations.

(i) Complete the missing labels on the diagram and name the technique.



Name of technique: Reflux [2]




(ii) Draw a labelled diagram to show apparatus set up for filtration under reduced pressure (vacuum filtration).




[2]




(b) This part is about amines.

(i) The table shows the structures and boiling points of three amines, which are structural isomers of C_3H_9N .

Amine	$CH_3CH_2CH_2NH_2$	$(CH_3)_2CHNH_2$	$(CH_3)_3N$
Skeletal formula			
Boiling point/ $^{\circ}C$	48–49 $^{\circ}C$	33–34 $^{\circ}C$	3–4 $^{\circ}C$

Explain the difference in the boiling points of the three amines.

 NH_2 no branches / longer chain, more points of contact / more surface in tetrahedral so stronger induced dipole-dipole interaction (London forces).
non-hindered N lone pairs

 NH_2 and  can form hydrogen bonds.  can't form H bonds

H bonds are stronger than London forces so more energy is needed to break H bonds. [4]

(ii) Amine A is a liquid at room temperature and pressure.

When vaporised, 0.202 g of the amine produces 72.0 cm³ of gas at 1.00 × 10⁵ Pa and 100 °C. The ¹³C NMR spectrum of amine A has 3 peaks.

Determine the molecular formula of A and suggest a possible structure for amine A.

$$n = \frac{PV}{RT}$$

$$R = 8.314$$


$$V = 72 \times 10^{-6} \text{ m}^3$$

$$T = 373 \text{ K}$$


$$P = 1 \times 10^5 \text{ Pa}$$

$$n = \frac{1 \times 10^5 \times 72 \times 10^{-6}}{8.314 \times 373} = 2.32 \times 10^{-3} \text{ mol}$$

$$\frac{0.202}{2.32 \times 10^{-3}} = 87$$



Molecular formula of A $\text{C}_5\text{H}_{13}\text{N}$




one example of possible structures

[6]

(c) The amino acid $\text{Z-H}_2\text{NCH=CHCOOH}$ can react to form a cyclic compound with the molecular formula $\text{C}_3\text{H}_3\text{NO}$ and one other product.

Complete the equation for this reaction.



look at what is missing from reactant

[2]

2* Transition metal ions can bond to ligands to form complex ions with different shapes.


Explain what is meant by the terms **ligand**, **coordination number** and **ligand substitution**, using suitable examples of **complex ions with different shapes**, limited to **monodentate ligands**.

Your answer should include diagrams and equations where appropriate. *each ligand [6] forms one coordinate bond*


ligand: Donates a lone pair to a metal ion forms dative covalent (coordinate) bond with metal ion

coordination number: number of coordinate bonds to metal ion

ligand substitution: one ligand replacing another





coordination number: 6 octahedral



Additional answer space if required


make sure O is bonding to central metal as O is donating its lone pair of electrons not H

coordination number: 4 tetrahedral



- 3 Propane, C_3H_8 , (boiling point $-42^\circ C$) is used as 'camping gas'. A student plans to determine the enthalpy change of combustion of propane, $\Delta_c H (C_3H_8)$, by two methods.

(a) The student first carries out an experiment using the apparatus below.



Results

Mass of propane burner before burning / g	99.218
Mass of propane burner after burning / g	98.976
Initial temperature / $^\circ C$	21.60
Maximum temperature reached / $^\circ C$	46.10

- (i) Determine the enthalpy change of combustion of propane, in $kJ mol^{-1}$.

Give your answer to 3 significant figures.

$$q = mc\Delta T$$

$$q = 100 \times 4.18 \times 24.5 = \pm 10241 \text{ J}$$

$$= \pm 10.241 \text{ kJ}$$

$$m = 100 \text{ g} = 100 \text{ cm}^3$$

$$c = 4.18 \text{ J/g}^\circ C$$

$$\Delta T = (46.1 - 21.6)$$

$$99.218 - 98.976 = 0.242 \text{ g of propane}$$

$$\frac{0.242}{44} = 0.0055 \text{ mol of propane}$$

$$\frac{10.241}{0.0055} = -1862 \text{ kJ mol}^{-1}$$

$\Delta_c H$ is exothermic

$$\frac{m \times c}{m \times R.F.M.}$$

$$(12 \times 3) + 8 = 44$$

$$\Delta_c H (C_3H_8) = -1860 \text{ kJ mol}^{-1} \text{ [3]}$$

(3sf.)

- (ii) The student finds that the experimental enthalpy change $\Delta_c H$ (C_3H_8) is much less exothermic than the accurate standard enthalpy change $\Delta_c H$ (C_3H_8) in databases.

One reason could be that the student's experiment had not been carried out under standard conditions.

Suggest **two** other reasons for this difference in enthalpy change.

1 heat loss/released to surroundings

2 incomplete combustion/reaction with oxygen/air/not enough burning


[1]

(3) Evaporation of water

- (b)* The student determines the standard enthalpy change of combustion of propane using the bond enthalpies in the table. An experiment is not needed.


Bond	Bond enthalpy /kJ mol ⁻¹
C-H	+413
C-C	+347
C=O	+805
O=O	+498
O-H	+464

The bond enthalpies can be used to determine the standard enthalpy change of reaction, $\Delta_r H$, for **equation 3.1**:



↑
treat as minus sign
Enthalpy change of vaporisation, $\Delta_{vap} H$

The standard enthalpy change of vaporisation of water, $\Delta_{vap} H$, is the enthalpy change for the conversion of 1 mol of $H_2O(l)$ into 1 mol of $H_2O(g)$ under standard conditions:



Determine the standard enthalpy change of combustion of propane (boiling point -42°C) using the $\Delta_r H$ value for **equation 3.1** and $\Delta_{\text{vap}} H$ for water. [6]

$$(2 \times 347) + (8 \times 413) + (5 \times 498) = \pm 6488 \text{ kJ mol}^{-1}$$



$$(6 \times 805) + (8 \times 464) = \pm 8542 \text{ kJ mol}^{-1}$$



$$6488 - 8542 = -2054 \text{ kJ mol}^{-1} = \Delta_r H$$

$$\Delta_r H - (4 \times \Delta_{\text{vap}} H)$$

→ 4 H₂O in equation 3.1

$$-2054 - (4 \times 40.65) = -2216.6 \text{ kJ mol}^{-1}$$

- 4 A student carries out an investigation to identify two metals, **M** and **X**, by two different methods.

- (a) The student is provided with a sample of metal **M**.

The student analyses metal **M** using a 'back-titration' technique:

- The metal is reacted with excess acid.
- The resulting solution is titrated to determine the amount of acid remaining after the reaction.

Stage 1

The student adds 100 cm^3 of 2.10 mol dm^{-3} HCl(aq) to 6.90 g of **M**.

An excess of HCl(aq) has been used to ensure that all of metal **M** reacts.

A redox reaction occurs, forming a solution containing **M** in the +2 oxidation state.


Stage 2

The resulting solution from **Stage 1** is made up to 250.0 cm^3 with distilled water.

Stage 3

A 25.00 cm^3 sample of the diluted solution from **Stage 2** is titrated with 0.320 mol dm^{-3} NaOH(aq) .

The NaOH(aq) reacts with excess HCl(aq) that remains in **Stage 1**:



The student repeats the titration to obtain concordant titres.

Titration results (The trial titre has been omitted,)


The burette readings have been recorded to the nearest 0.05 cm^3 .

	1	2	3
Final reading/ cm^3	27.80	37.55	32.20
Initial reading/ cm^3	0.50	10.00	5.00
titre/ cm^3	<u>27.30</u>	<u>27.55</u>	<u>27.20</u>

concordant

- (i) In **Stage 1**, a redox reaction takes place between **M** and HCl(aq) , forming hydrogen and a solution containing **M** in the +2 oxidation state.


Write an overall equation, with state symbols, for this reaction. Write half-equations for the oxidation and reduction processes.



(increase oxidation state)



(decrease oxidation state)



[3]

(ii) In Stage 1, suggest two observations that would confirm that all of metal M has reacted.

1. bubbles/effervescence stops

2. M/metal disappears

[2]

(iii) In Stage 3, write the ionic equation for the reaction taking place in the titration.



(iv) Metal M can be identified following the steps below.

- The amount, in mol, of excess HCl(aq) that remains after the reaction of M with HCl(aq).
- The amount, in mol, of HCl(aq) that reacted with M.
- The identity of metal M.

Analyse the results to identify metal M.

$$\frac{27.30 + 27.20}{2} = 27.25 \text{ cm}^3 \text{ mean titre}$$



$$27.25 \times 10^{-3} \times 0.320 = 8.72 \times 10^{-3} \text{ mol in } 25 \text{ cm}^3 \text{ (HCl)}$$

$$8.72 \times 10^{-3} \times 10 = 8.72 \times 10^{-2} \text{ mol in } 250 \text{ cm}^3 \text{ (HCl)}$$

$$0.210 - 8.72 \times 10^{-2} = 0.1228 \text{ mol HCl reacted with M}$$

$$\frac{0.1228}{2} = 0.0614 \text{ mol of M reacted}$$


$$\frac{6.90}{0.0614} = 112.4 = \text{Cd}$$



Metal M = Cd [6]

- (b) The student is provided with the carbonate of an unknown metal, X_2CO_3 .

The student measures the mass loss when the X_2CO_3 is reacted with an excess of hydrochloric acid. The equation is shown below.



The reaction is carried out using this method:

- Step 1** Add 100 cm³ HCl(aq) to a conical flask and weigh.
Step 2 Add X_2CO_3 to the conical flask and immediately reweigh.
Step 3 After 5 minutes, reweigh the conical flask and contents.

Results

Mass of conical flask + HCl(aq)	172.93g
Mass of conical flask + X_2CO_3 + HCl(aq) before reaction	187.50g
Mass of conical flask + contents after 5 minutes	184.75g

- (i) Calculate the amount, in mol, of CO_2 released in the reaction.

$$187.50 - 184.75 = 2.75 \text{ g of } CO_2$$

$$\frac{2.75}{12 + (16 \times 2)} = 0.0625$$

Amount of CO_2 = 0.0625 mol [1]



- (ii) Calculate the molar mass of X_2CO_3 and identify metal X.

$$0.0625 \text{ mol of } X_2CO_3$$

$$187.50 - 172.93 = 14.57 \text{ g of } X_2CO_3$$

$$\frac{14.57}{0.0625} = 233.12 \text{ g mol}^{-1}$$

$$X = Rb$$

$$\frac{233.12 - (12 + (16 \times 3))}{2} = 86.56 = Rb$$

Molar mass of X_2CO_3 = 233.12 g mol⁻¹ Metal X = Rb [3]

- (c) After analysing the results, the student was told that their molar mass of X_2CO_3 was incorrect.

The student evaluated the experiment for possible reasons for the incorrect result.

- (i) The student wondered whether the reaction was complete when the mass was recorded after 5 minutes (Step 3).

How could the student modify the experimental procedure to be confident that the reaction was complete?

reweigh to constant mass.....

.....

..... [1]

- (ii) The student finds out that carbon dioxide is slightly soluble in water.

State and explain how the solubility of CO_2 would affect the calculated molar mass of X_2CO_3 .

• mass (CO_2) loss would be.....

smaller/ mass X_2CO_3 reacted seems
to be less.....

• molar mass would be greater..... [2]

- 5 The equilibrium constant K_p and temperature T (in K) are linked by the mathematical relationship shown in equation 5.1 (R = Gas constant in $\text{J mol}^{-1} \text{K}^{-1}$ and ΔH is enthalpy change in J mol^{-1}).

$$\ln K_p = -\frac{\Delta H}{R} \times \frac{1}{T} + \frac{\Delta S}{R} \quad \text{Equation 5.1}$$

$$y = m \times x + c$$

- (a) The table shows the values of K_p at different temperatures for an equilibrium.

Complete the table by adding the missing values of $\frac{1}{T}$ and $\ln K_p$.

$$K_p = \frac{p(\text{products})}{p(\text{reactants})}$$

Temperature, T/K	400	500	600	700	800
K_p	3.00×10^{58}	5.86×10^{45}	1.83×10^{37}	1.46×10^{31}	1.14×10^{26}
$\frac{1}{T}/\text{K}^{-1}$	2.50×10^{-3}	2.00×10^{-3}	1.67×10^{-3}	1.43×10^{-3}	1.25×10^{-3}
$\ln K_p$	135	105	86	72	60

[2]

- (b) State and explain how increasing the temperature affects the position of this equilibrium and whether the forward reaction is exothermic or endothermic.

Equilibrium position shifts to the left. Forward reaction is exothermic.

As temperature increased the equilibrium position moves in the endothermic direction which in this case is the backwards reaction.

- (c) Plot a graph of $\ln K_p$ against $\frac{1}{T}$ using the axes provided on the opposite page.

Use your graph and equation 5.1 to determine ΔH , in kJ mol^{-1} , for this equilibrium.

Give your answer to 3 significant figures.

$$\frac{\Delta y}{\Delta x} = 57142.85714 \quad \frac{\Delta y}{\Delta x} = \frac{-\Delta H}{R}$$

$$-\Delta H = 475085.7143 \text{ J mol}^{-1}$$

$$\Delta H = -475.085 \text{ kJ mol}^{-1}$$


$$\Delta H = -475 \text{ kJ mol}^{-1} \quad [4]$$

- (d) Explain how ΔS could be calculated from a graph of $\ln K_p$ against $\frac{1}{T}$.

$$y\text{-intercept} = \frac{\Delta S}{R}$$

$$\Delta S = R \times y\text{-intercept}$$

[2]



- 6 This question is about two different types of acid found in organic compounds, carboxylic acids and sulfonic acids, as shown in Fig. 6.1.




Fig. 6.1

- (a) Complete Table 6.1 to predict bond angles *a* and *b* and name the shapes which makes these bond angles in the functional groups of carboxylic acids and sulfonic acids.



Type of acid	Acid	Bond angle	Name of shape
Carboxylic acid		120°	trigonal planar
Sulfonic acid		109.5°	non-linear

Table 6.1

lone pairs repel more than bonded pairs

[2]

- (b) Ethanoic acid, CH_3COOH , and methanesulfonic acid, $\text{CH}_3\text{SO}_2\text{OH}$, are both monobasic acids. The $\text{p}K_a$ values are shown in the table.

Acid		$\text{p}K_a$
Ethanoic acid	CH_3COOH	4.76
Methanesulfonic acid	$\text{CH}_3\text{SO}_2\text{OH}$	-1.90

$\text{p}K_a = \text{pH}$

A student suggests that $1.0 \text{ mol dm}^{-3} \text{ CH}_3\text{SO}_2\text{OH}$ should have a lower pH value than $1.0 \text{ mol dm}^{-3} \text{ CH}_3\text{COOH}$.

Write an equation, showing conjugate acid-base pairs, for the equilibrium of $\text{CH}_3\text{SO}_2\text{OH}$ with water and explain, with reasons, whether the student is correct.

Label the conjugate acid-base pairs: **A1, B1** and **A2, B2**.



acids: proton donors

bases: proton acceptors

$\text{CH}_3\text{SO}_2\text{OH}$ is a stronger acid / dissociates more

student is correct $\text{CH}_3\text{SO}_2\text{OH}$ has a lower [4]
 $\text{p}K_a / \text{pH}$ / higher K_a / $[\text{H}^+]$


(c) Carboxylic acids and sulfonic acids both form esters.

Sulfonic acid esters can be hydrolysed by aqueous alkali.
The equation shows the alkaline hydrolysis of a sulfonic acid ester.



In the 3 boxes below, add curly arrows to show the mechanism for this reaction.

In the first box, the hydroxide ion acts as a nucleophile.



look at what you want to make at each stage and work backwards to draw curly arrows

[4]

END OF QUESTION PAPER